AD	•	

Award Number: W81XWH-04-1-0612

TITLE: Identification of Genes Involved in Breast Tumor Invasion Utilizing a Ubiquitin-Mediated Proteolysis In Vitro Screen

PRINCIPAL INVESTIGATOR: Charles H. Spruck, Ph.D.

CONTRACTING ORGANIZATION: Sidney Kimmel Cancer Center

San Diego, CA 92121

REPORT DATE: October 2005

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20060503022

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188			
Public reporting burden for this of	collection of information is estima	ited to average 1 hour per respo	nse, including the time for review	ing instructions, search	ing existing data sources, gathering and maintaining the		
this burden to Department of De 4302. Respondents should be	fense, Washington Headquarter aware that notwithstanding any c	s Services, Directorate for Inform	nation Operations and Reports (C shall be subject to any penalty fo	704-0188), 1215 Jeffer:	ection of information, including suggestions for reducing son Davis Highway, Suite 1204, Arlington, VA 22202- a collection of information if it does not display a currently		
1. REPORT DATE		REPORT TYPE			ATES COVERED		
01-10-2005		nnual		30	Sep 2004 –29 Sep2005		
4. TITLE AND SUBTITE	E			5a. C	CONTRACT NUMBER		
Identification of Ger	nes Involved in Brea	ast Tumor Invasion	Utilizing a Ubiquitin-				
-Mediated Proteolysis In Vitro Screen					GRANT NUMBER		
					1XWH-04- Identification of		
				5c. F	PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. F	PROJECT NUMBER		
Charles H. Spruck, Ph.D.				5e. 1	ASK NUMBER		
					WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)					ERFORMING ORGANIZATION REPORT		
Sidney Kimmel Cancer Center				N	UMBER		
San Diego, CA 92	121						
		•					
0 SDONGODING (1)	NITODING ACTUOY	ME(C) AND ADDECC	(/ES)		EDONGOD/MONITORIE ACRONICATO		
U.S. Army Medical		AME(S) AND ADDRESS eriel Command	n(EO)	10.8	SPONSOR/MONITOR'S ACRONYM(S)		
Fort Detrick, Maryla							
				11.5	SPONSOR/MONITOR'S REPORT		
				l l	NUMBER(S)		
12. DISTRIBUTION / A							
Approved for Public Release; Distribution Unlimited							
40 OURDI SEESENS	(NOTE)						
13. SUPPLEMENTARY	NOTES						
•							
14. ABSTRACT	e evalore the notest	ial use of ubiquities	denendent protocky	sie ae a "rover	ea ganatics" tool in functional		
					se genetics" tool in functional		
yenomics studies.	vve wiii uevelop a l	enoviral-based syst	tem that artificially to Looptide librory	aryets random	cellular proteins to the proteolytic		
machinery for degradation. To achieve this, a randomized peptide library will be linked to a segment of the F-box motif of beta-							
TrCP, the F-box protein that mediates the ubiquitination of I(kappa)B(alpha) and (beta)-catenin via the multimeric SCF							
ubiquitin ligase. The resultant chimeric proteins are expected to direct any interacting proteins that are otherwise stable, to the							
SCF ligase for ubiquitination. As proof of principle, we will use this system in a loss-of-function <i>in vitro</i> assay to identify							
putative genes involved in breast tumor invasion. MDA-MB-231 breast tumor-derived cells will be tranduced with the retroviral							
chimera library and peptides that confer the ability to invade through an artificial extracellular matrix will be isolated using a							
modified Boyden chamber assay. Following multiple rounds of selection, the chimeras that provide invasion properties will be							
confirmed using tumorgenicity assays in nude mice. If successful, ubiquitination-based functional assays will undoubtedly							
contribute to the identification of potential protein targets for therapeutic intervention in breast cancer.							
AL OUD FOT TEDMO							
15. SUBJECT TERMS Invasion, Ubiquitin, Proteolysis, SCF, F-box, Functional Genomics							
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC		
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area		
Ŭ	Ü	Ŭ	שט	5	code)		

Table of Contents

Cover	1
SF 298	2
Table of Contents	3
Introduction	4
Body	4-5
Key Research Accomplishments	5
Reportable Outcomes	5
Conclusions	5
References	5
Appendices	5

A. Introduction

The goal of this research project was to establish a functional genomics screen that utilizes a cell's own ubiquitin-dependent proteolysis system to screen for proteins involved in breast tumor invasion. The experimental design of the project was to express a randomized peptide library attached to a segment of the F-box domain of the protein β-TrCP in MDA-MB-231 breast tumor cells. The chimeric proteins are expected to bind to and recruit otherwise stable cellular proteins to the SCF ubiquitin ligase for ubiquitin-dependent degradation. Cells that express this construct would then be selected for a gain of invasion function using a Boyden chamber assay.

B. Body

We have asked for and were granted a one-year extension for the completion of this research project from the DOD Breast Cancer Research Program. The extension was requested based on unexpected complications in constructing and amplifying the proteolysis targeting vector library of sufficient molecular complexity for functional genomics experiments. This problem was due to an inefficient transformation efficiency of *E. coli* cells which has since been overcome (described below).

We have amplified a segment of the protein β -TrCP pertaining to the F-box domain from expression plasmid pFlag-CMV2- β -TrCP using the Advantage HF2 high fidelity polymerase (Clontech). We designed the oligonucleotides to include a 5' Flag epitope tag and 5' and 3' cloning sites for directional cloning. We sequenced the amplified DNA segment and found that it to be free of mutations (completion of point #1 of SOW).

We then devised an experimental strategy to generate the DNA segment that encodes the random peptide library. This strategy was based on annealing a degenerate oligonucleotide with a complementary anchor oligonucleotide (both synthesized by Sigma-Genosys). The oligonucleotides were designed to contain 5' and 3' restriction endonuclease sites for directional cloning, sequence which encodes a 5' poly-glycine linker sequence, and a degenerate (N)²¹ DNA sequence that encodes peptides of random 7 amino acids. The sequence content of the degenerate oligonucleotide was designed to minimize the possibility of the introduction of a stop codon in the peptide library. We annealed the degenerate and anchor oligonucleotides and constructed the double-stranded DNA peptide library cassette by incubation with Klenow polymerase (Clontech). The efficiency of the annealing and synthesis reactions was evaluated by agarose gel electrophoresis (completion of SOW point #2).

We constructed the retroviral-based proteolysis targeting vector by first cloning the β -TrCP F-box segment synthesized above into the retroviral vector pLPC (a previous gift from S. Lowe, Cold Spring Harbor Laboratory). This vector was chosen due to its high level of expression from a CMV minimal promoter. We then directionally cloned the random peptide library cassette synthesized above in the cloning site 3' of the β -TrCP cassette. The efficiency of the ligation reaction was evaluated by agarose gel electrophoresis (completion of point #3 of SOW)

In order to perform a comprehensive functional genetic screen for genes involved in breast tumor invasion, we estimated that a molecular complexity of $>10^{10}$ different proteolysis targeting plasmids would be required. However, we encountered a "bottleneck" problem in the amplification of the ligated proteolysis targeting vector. Using various commercially available strains of chemically competent E. coli, we could not

achieve a transformation frequency $>10^7/\mu g$ for the proteolysis targeting construct. Over the last 2-3 months, we have compared various commercially available chemically and electrocompetent strains of E.~coli to optimize the transformation efficiency. We have now optimized our transformation procedure using a commercially available electrocompetent strain of E.~coli that provides a transformation efficiency $>10^{10}/\mu g$ proteolysis targeting vector. This transformation efficiency avoids the "bottle-neck" on molecular complexity encountered in proteolysis targeting vector amplification and will allow for us complete the remaining points outlined in the SOW in the coming year.

C. Key research Accomplishments

Construction of a retroviral-based proteolysis targeting vector for expression of 7mer random peptides linked to a β -TrCP F-box domain.

D. Reportable Outcomes

<u>Abstract:</u> Tan, YM, Spruck, C. Identification of genes involved in breast tumor invasion utilizing an ubiquitin-mediated proteolysis in vitro screen. Era of Hope meeting 2005, sponsored by the DOD Breast Cancer Research Program

E. Conclusions

A one-year extension was asked for and granted from the DOD Breast Cancer Research Program for the completion of this research. The estimated completion date of the project is November 2006.

G. References

None

H. Appendices

None